

**CAG Survey of the use of Systemic Non-opioid Adjuvant Analgesics (NOAA), excluding Paracetamol and NSAIDs, for Colorectal Surgery:**

**NOAAs – Gabapentoids, Alpha2 agonists, Ketamine, Lidocaine, Magnesium, Dexamethasone**

Please circle or tick (in the table) appropriately. Please complete entire survey form. If any comment please write appropriately. Many thanks.

- **How frequently do you anaesthetise colorectal surgical patients? -**

Regularly (weekly/biweekly)                      Monthly                      Occasionally                      N/A

- **Do use NOAA for?-**

Elective cases                      Emergency cases                      Both                      Do not use

- **Do you use NOAA for?–**

Open surgery                      Laparoscopic surgery                      Both                      Do not use

- **Do you use NOAA for colorectal surgery? –**

Routinely                      Randomly                      Never

- **Does NOAA has role for enhanced recovery for colorectal surgical patients?-**

Yes                      No                      Do not know

- **Is NOAA a part of guideline for enhanced recovery for colorectal surgery in your hospital? -**

Yes                      No                      Do not know

- **Is there a guideline for use of NOAA In your department/hospital ?–**

Yes                      No                      Do not know

- **Commonly I use following NOAA for colorectal surgical patients (Please tick & mention the dose -**

	Preop /Dose	Intraoperative /Dose	Post op Day 1/Dose	Post op day 2/Dose
Gabapentin				
Pregabalin				
Clonidine				
Dexmedetomidine				
Ketamine				
Lidocaine infusion				
Magnesium				
Dexamethasone				
Other (Please mention the name)-				



